

A GUIDE TO YOUR KNEE REPLACEMENT EXPERIENCE



This guide will assist in preparing you for your upcoming knee replacement procedure at UnaSource Surgery Center. Being informed and prepared will be a large factor in your surgery's success.

Please review this booklet, share it with the friends and family members who will be assisting you during recovery, and bring it with you to all events related to your surgery.

UnaSource
Surgery Center

Table of Contents

Introduction..... 1-2

Joint Replacement Risks

 General Risks.....5-6

 Complications specific to procedure.....6

Preparing for your Outpatient Total Joint Replacement

 Timeline.....9

 Registering for Surgery.....10

 Online Joint Replacement Education Video.....10

 Medical Clearance & Medications.....11

 Home Preparation Checklist.....12

What to Expect on your Day of Surgery

 Day Before Surgery.....15

 Your Surgical Experience.....15

 Discharge.....16

What to Expect During Recovery

 Pain Management.....19

 Home Health & Physical Therapy.....20

 Moving Around & Precautions.....20

 Caring for your Incision.....21

 When to Call the Surgeon.....21

 Medication Tracker.....22-23

 Knee Replacement Pre-Op Instructions Checklist.....24

 Notes.....25-27



Joint replacement is a surgical procedure in which the worn, damaged surfaces of the joint are removed and replaced with new artificial parts. Your doctor may consider joint replacement if you have severe pain which limits daily activities and is not relieved with medications, injections, physical therapy, or other treatments. The goal of the surgery is to relieve pain and restore the alignment and function of your joint.

Joint replacement provides the most dramatic relief of arthritis pain available to patients.

Traditionally, joint replacement has been an inpatient surgical procedure. With the introduction of minimally invasive techniques, modern pain relief protocols, improved anesthetic techniques, and rapid rehabilitation, surgeons can perform total joint replacement procedures on an outpatient basis.

OUTPATIENT JOINT REPLACEMENT IS THE SAME AS TRADITIONAL JOINT REPLACEMENT BUT THE PATIENT CAN GO HOME ON THE SAME DAY OF SURGERY. THIS ALLOWS PATIENTS TO RECOVER IN THE COMFORT OF THEIR OWN HOME INSTEAD OF OVERNIGHT IN A HOSPITAL.



With outpatient joint replacement, patients can often return to normal activities quicker and with less pain than with traditional joint replacement surgery.

Other advantages of outpatient joint replacement include:

- Rapid recovery
- Decreased hospital stay
- Improved pain management protocols
- Early mobilization
- Careful home monitoring
- Fewer complications
- Improved outcomes
- Increased patient satisfaction

Knee replacements done at UnaSource:



Uni-Compartmental “Partial” Knee Arthroplasty

Either the medial or lateral portion of the femur (thigh bone), tibia (shin bone) and kneecap are resurfaced



Total Knee Arthroplasty

Both the medial and lateral portions of the femur (thigh bone), tibia (shin bone) and kneecap are resurfaced



KNEE REPLACEMENT RISKS

As with any major surgery, there are a number of potential complications related to joint replacement surgery. The general risks of joint replacement surgery include: heart attack, stroke, infection, pneumonia, bleeding, blood clot, urinary dysfunction, and nerve or blood vessel damage. This list is not all-inclusive.

Infection

To reduce the risk of a surgical site infection, your nose will be swabbed with betadine swabs. You will also shower with Chlorhexidine Gluconate 4% (CHG) skin cleanser for 2 days before surgery and on the morning of your surgery to reduce the bacteria on your skin (see page 11). On the day of surgery at UnaSource Surgery Center you will receive an antibiotic.

To reduce the risk of a surgical site infection:

- ★ Practice good hygiene during your recovery
- ★ Wash your hands
- ★ Wear clean clothes
- ★ Use clean linens and bedding
- ★ Avoid touching the bandage on your wound

Pneumonia

To reduce the risk of pneumonia:

- ★ Stop smoking now. Smoking increases your anesthesia risks, increases your risk of pneumonia after surgery, and slows tissue repair at the site of your surgery.
- ★ Deep breathe. After your surgery during your recovery, take at least 10 deep focused breaths every hour while you are awake. Try to do this as long as you are taking pain medication.
- ★ Avoid others who are sick, especially those with a cough.

Bleeding

To reduce the risk of bleeding:

- ★ Stop taking anti-inflammatory medications 7 days prior to surgery, or as directed by your surgeon. Examples of these medications are: Motrin, Aleve, Mobic, Naprosyn, aspirin, etc.
- ★ If you are on blood thinners such as Coumadin (warfarin), Plavix, Xarelto, Eliquis, Lovenox or aspirin – you may need to stop taking these medications prior to surgery. Talk to your surgeon and prescribing provider to determine what is best for you.

Blood Clot

Blood clots commonly form in the legs. A blood clot that lodges in the veins of the leg is called a deep vein thrombosis (DVT). To reduce your risk of a DVT, your surgeon will prescribe a blood thinner such as aspirin for you to take after your surgery. You may be prescribed a sequential compression device (SCD) and wraps to wear on both legs; the SCD will be worn for 2 weeks during times when you are resting at home, or as directed.



To reduce the risk of a blood clot:

- ★ Use the prescribed sequential compression device as instructed.
- ★ Get up and walk; moving around keeps your blood circulating.
- ★ Perform ankle pumps and rotate your ankles clockwise and counter-clockwise every hour while at rest.
- ★ Contract your thigh muscles every hour you are awake.

Urinary or Bladder Issues

After surgery it can be difficult to urinate and your stream may be weak. You will be required to urinate prior to discharge. You should be able to urinate within 6 hours after your surgery. Turning on a faucet may help bring on the urge to urinate. Notify your surgeon if you are not able to urinate within this timeframe.

Nerve Damage

All patients will notice numbness in the skin around the incision site. The area of numbness usually decreases in size over time.

Knee Replacement Complications

Stiffness: Difficulty with full extension (straightening) or full flexion (bending) of the knee. Sometimes due to the design of the knee implant. The most important predictor of postoperative knee motion is preoperative knee motion.

Instability: Very rare loose sensation in the knee. May be treated over time with bracing and strengthening exercises.



PREPARING FOR OUTPATIENT KNEE REPLACEMENT

UnaSource Surgery Center Knee Replacement Experience Timeline*

*your individual experience may vary

2-4 Weeks Before Surgery	Watch the Joint Replacement Education video on unasourcesurgery.com
	Stop taking certain medications as instructed.
	UnaSource Surgery Center Patient Registration will call to verify your demographic and insurance information.
	Look for an email from our RN Ortho Coordinators with steps to take prior to surgery.
1-2 Weeks Before Surgery	A UnaSource Ortho Coordinator will call to go over your health history and answer questions. Be sure to have a complete list of your medications available for the call.
2 Days Before Surgery	Start showering with Chlorhexidine Gluconate.
Day Before Surgery	Shower with Chlorhexidine Gluconate.
	UnaSource Surgery Center will call between 3-5pm with your surgery arrival time.
	Wash bedding and towels.
	Do not eat after midnight.
Day of Surgery	Shower with Chlorhexidine Gluconate at home prior to surgery.
	Replace bedding with fresh linens prior to surgery.
	Take the medications you have been instructed to take with only a sip of water.
Day After Surgery	Your home care company will call to set up home visits if not previously arranged.
	A UnaSource Surgery Center Ortho Coordinator will call to follow up.
	Call your surgeon's office to schedule follow-up appointment.
48 Hrs After Surgery	A UnaSource Surgery Center Ortho Coordinator will call to follow up.
4 Days After Surgery	Remove pain pump catheter, if a pain pump had been used.
7 Days After Surgery	A UnaSource Surgery Center Ortho Coordinator will call to follow up.
30/60/90 Days After Surgery	You will receive a follow up email from UnaSource.



Pre Surgical Health Assessment

Once you are scheduled for joint replacement at UnaSource Surgery Center, you need to complete a pre-surgical health assessment for the anesthesiologists who will be caring for you on the day of surgery. An Ortho Coordinator will call you 1-2 weeks prior to surgery to complete your health assessment. Be sure to have a complete list of your medications available for the call.

Joint Replacement Education Video

You and any support person or family member who will be helping you recover at home after surgery should watch our online Joint Replacement Education video. The video will cover:

- How to prepare for your surgery
- What to expect on your day of surgery
- What your recovery plan at home will involve

There is no surgical/graphic footage in this video.

Enter the patient's name and email at: www.unasourcesurgery.com/joint-replacement to watch.

Use this section to write down the questions you'd like to ask our team after watching the video.

Medical Clearance

Your surgeon may require a medical or cardiac evaluation prior to surgery. You should see your primary care doctor and/or cardiologist if this is required. The results of the medical evaluations need to be sent to your surgeon's office.

Dental Work

If you have not seen a dentist in the past 6 months, make an appointment as soon as possible for a dental exam. You should not have any major dental work or a cleaning in the month prior to your joint replacement surgery or for 3 months after your surgery.

Flu Shot/Covid Shot

If you choose to get one of these vaccinations, please make sure it is completed at least 4 weeks prior to the surgery date.

Medications

Stop taking the following medications 7-14 days prior to surgery, or as directed:

- Anti-inflammatory (Motrin, Aleve, Mobic, Naprosyn, aspirin, etc)
- Vitamins and Supplements (fish oil, calcium, glucosamine)
- Rheumatologic (Enbrel, Remicade, Methotrexate)

If you are on blood thinners such as Coumadin (Warfarin), Plavix, Xarelto, Eliquis, Lovenox or aspirin – you may need to stop taking these medications prior to surgery. Continue taking pain medication and narcotics, as well as blood pressure, diabetic, and heart medications.

Betadine Swab

Your nose will be swabbed with a betadine swab the day of surgery to eliminate any Staphylococcus Aureus bacteria that may be present. Thirty-five percent of the population carry this bacteria in their nose, which can transfer to your skin. Studies have shown that eliminating this bacteria from your nose prior to surgery reduces the risk of a surgical site infection.

Chlorhexidine Gluconate 4% (CHG) Skin Cleanser

Showering with this soap has been shown to decrease the bacteria level on your skin which may reduce your risk of a surgical site infection. Plan to purchase the CHG skin cleanser and shower with it the 2 days prior to your surgery and on the morning of your surgery before you leave your home for the Center.



Preparing for Recovery at Home

You must have a support person who will accompany you at UnaSource Surgery Center on the day of your surgery and stay with you for the first few days after surgery. It is ideal that your support person also watches our online Joint Replacement video.

You will be going home with a walker. You may consider attaching a basket or bag to the walker as it is not safe to carry items when using it.

Make sure pathways between rooms are clear:

- ☐ Remove doormats and throw rugs
- ☐ Secure floor-level cords
- ☐ Remove clutter from the floors



Pets can be a trip hazard too!

Complete shopping, housekeeping and laundry prior to your surgery date:

- ☐ Put newly laundered linens on your bed the morning of your surgery and ensure you have a supply of clean bath towels – using clean linens and towels may reduce the risk of getting a surgical site infection
- ☐ Move items you will need or frequently use in the kitchen, cupboards & bathroom to waist level for easy access
- ☐ Get a shower chair
- ☐ Get a raised toilet seat
- ☐ Prepare for meals and shop in advance



WHAT TO EXPECT ON YOUR DAY OF SURGERY

Day Before Surgery

UnaSource Surgery Center will call between 3-5pm the day before your surgery to notify you of your arrival time. For those having surgery on a Monday, this call will be made on the Friday prior to your surgery date. You are typically asked to arrive at UnaSource Surgery Center 2 hours prior to your scheduled surgery time. During this phone call, the UnaSource team member will review medication instructions, let you know when to stop eating and drinking, and answer any questions you may have.

Your Surgical Experience

On your day of surgery, the support person accompanying you to UnaSource Surgery Center must be a responsible adult. Wear loose, comfortable clothes and non-skid, sturdy shoes. Do not wear jewelry or contact lenses. Remember to bring:

- Driver's License
- Insurance Cards
- Any durable medical equipment prescribed by your surgeon

Medications to take before surgery	Medications to stop taking before surgery

After you have registered at the Front Desk, you will be escorted to a private Patient Care room; the person accompanying you will join you after you are admitted. You will meet the anesthesiologist who will talk about the type of anesthesia that will be used during your surgery. Before surgery, the anesthesiologist may provide a peripheral nerve block or insert a pain pump catheter to keep you comfortable after surgery. Also before surgery, you will receive a sequential compression device with stockings that pump air around your calves.

During your surgery, the responsible adult accompanying you at UnaSource will wait in our Lobby. Beverages, Wi-Fi, reading materials, television and comfortable seating are available for your waiting guest.

After surgery you will return to your private Patient Care room for recovery and your surgeon will speak with the person accompanying you in our private consultation room.

If your surgeon sent your prescriptions to the pharmacy on the UnaSource campus, a nurse will let your support person know to pick them up while you are in surgery.

Once you wake up you will be offered something to drink and a light snack.

Your Team of Professionals

Surgeon	Operating Room Nurse
Physician Assistant	Surgical Technologist
Anesthesiologist	Recovery Nurse
CRNA (Nurse Anesthetist)	Clinical Technician/Nursing Assistant
Pre-Operative Nurse	Physical Therapist

Discharge Planning

A registered nurse will review all of your surgeon's orders and instructions with you and the person accompanying you. You will receive a folder to take home with copies of all the orders and instructions.

If you have a pain pump, the nurse will provide instructions on how to use the pain pump and also how to remove the small catheter at home once the pain pump is empty. The pain pump provides a continuous flow of local anesthesia (numbing medicine) around the nerves close to the surgical site. The pain pump will last for 4 days.

A physical therapist will review the use of your walker, how to use the stairs with your walker, and will also review the exercises ordered by your surgeon. You may put weight on your leg right away. The walker will be used until your surgeon advances you to the cane.

You will have a knee brace on and will need to wear the brace until the peripheral nerve block has worn off or the pain pump catheter is removed. **This knee brace is to help prevent you from falling.**

You may be sent home with a sequential compression device to wear at home for 2 weeks at times when you are inactive, or as directed. Calf pain, tenderness, redness, or swelling must be reported to your surgeon immediately.


Your surgeon will prescribe the appropriate post op meds for you.



WHAT TO EXPECT DURING RECOVERY

Pain Management

Most people will experience discomfort after surgery and it is normal. The pain scale below will be used to determine your pain level during your recovery at UnaSource Surgery Center and also once you are at home.



The goal of pain management is to lessen discomfort so you can rest, be mobile, and effectively participate in physical therapy. The plan to manage your postoperative pain starts before you even have surgery.

Understanding Your Pain Management Options

On the day of your surgery, you will meet with your anesthesia and surgical team to finalize your pain management plan. A multimodal approach is used to keep you comfortable. The plan will include options to help numb the knee after surgery.

Your surgeon and anesthesia team will determine which option is the best for you.

Common options include:

- **Local nerve block:** This is a long-acting nerve block that numbs the nerves that supply the area around the knee and helps provide extended pain relief after surgery.
- **Pain Pump:** This is a small catheter that delivers numbing medication continuously to the knee area and helps provide ongoing pain relief after surgery.
- **Local Injections:** Numbing medication may be injected around the surgical area by the surgeon during the procedure, helping with pain relief after surgery.

These methods work together to minimize your post-surgical discomfort and help you recover more comfortably.

Important to note: If you receive a nerve block or pain pump, your leg may feel partially numb or weak for some time after surgery. You may not feel discomfort from pressure or injury. It is important to wear the knee immobilizer when walking as directed and follow all precautions to protect your knee and keep you safe during this time. This numbness is temporary and will improve as the numbing medication wears off.

At home, you should take the pain medication on a regular schedule and as prescribed so you stay ahead of the pain. It can take some time to get the pain to a tolerable level if the interval between doses is too long. Take pain medication with food to reduce the chance of becoming nauseated.

Placing ice over your surgical site and keeping your surgical site elevated may also lessen discomfort. Use an ice pack for 20-30 minutes 4-6 times per day. Do not place the ice pack directly on skin.

Side Effects from Pain Medications

Nausea: Take pain medications with food to reduce the chance of nausea.

Constipation: Common while taking pain medications. A stool softener such as Colace will be prescribed by your surgeon to minimize constipation. A high fiber diet including prune juice, and increased activity will also help lessen constipation.

Home Health Care and Physical Therapy

- In home Physical Therapy may be arranged for you based on your surgeon's preference.
- Outpatient Physical Therapy should be arranged per your surgeon's directive.

Moving Around

The peripheral nerve block you received from the anesthesiologist will cause numbness and possible weakness to your surgical leg.

To help prevent falling, it is important that you are cautious when moving around and that you use your walker at all times.

- Wear non-skid shoes
- Keep the knee brace on every time you are up walking
- Ambulate with the walker at all times

Plan on using the walker for 1 – 3 weeks after surgery and then a cane for the next 6 – 8 weeks.

Remember to use the sequential compression device whenever you are not up and walking. Do NOT walk with the sequential device wraps on your calves - they will slip down to your ankles and create a fall hazard.

Knee Replacement Precautions

- ✓ Elevate your surgical leg when resting.
- ✓ Use a shower chair while showering.
- ✗ Do not place a pillow under or behind your knee. Instead, place the pillow under your ankle when elevating.
- ✗ Do not kneel on surgical knee.



Caring for your Incision

- Always wash your hands before and after touching the incision area.
- Wear clean clothing every day and use clean linens.
- If you have an ace wrap around your knee, you may remove it that first evening at home.
- The dressing over your incision should stay on for 7 days, or as directed by your surgeon. The dressing can get wet in the shower.
- You may shower 24 hours after your surgery, unless you have a pain pump. If you do have a pain pump you may shower when the pain pump is removed, on the fourth day after surgery.
- When showering after the dressing is removed, you may gently wash over the incision with a mild soap and water.
- Do not bathe in a tub for 2 weeks after surgery, or as directed by your surgeon.
- Do not apply any ointments or creams to your incision unless your surgeon has asked you to do so.
- Some drainage from the surgical site can be expected during the first 7 days after your surgery.

Some swelling and discoloration over or around the incision is normal and usually noticed towards the end of the day. Swelling and bruising may move down the leg and even into the feet.

To minimize swelling:

- Elevate the operative leg on pillows several times a day for 30 minutes each time. Your toes should be above the level of your heart.
- Place ice over the incision and surrounding areas as a mild pain reliever.
- Do not use heat as it may cause more swelling.

When to Call the Surgeon's Office



- A temperature above 100.5 F taken at least twice over four hours
- Appearance of red, hot, swollen incision
- A sudden increase or persistent increase in swelling in your surgical leg
- Unusual pain in your calf, knee, thigh or hip
- A calf that becomes swollen, tender, warm or reddened
- New or an increase in drainage from your incision

**IF YOU HAVE SHORTNESS OF BREATH OR CHEST PAIN,
CALL 911 IMMEDIATELY.**

Post-Op Prescription Tracker

Taking the medications as your surgeon has prescribed is an important part of a successful recovery. We recognize that keeping all your medications straight can be a challenge and have created this tool to assist you. Be sure to follow your surgeon's orders in the event you receive conflicting information.

Instructions:

Add the names of the prescribed medication in the empty light blue or tan box at the top of each section of the table. Then complete the dosage, frequency, and reason for taking that medicine (ie "Nausea") as written on your pill bottle. The boxes below the clock images are to record the time you took the medication.

Example:

Medicine	Medication Name				Medication Name			
Rx	# pill(s) every # hours as needed for <i>reason</i>				# pill(s) every # hours as needed for <i>reason</i>			
DATE								
Date Taken	Time 1st dose taken	Time 2nd dose taken			Time 1st dose taken	Time 2nd dose taken		

Medicine								
Rx	___ pill(s) every ___ hours as needed for _____				___ pill(s) every ___ hours as needed for _____			
DATE								

Medicine								
Rx	___ pill(s) every ___ hours as needed for _____				___ pill(s) every ___ hours as needed for _____			
DATE								

Medicine								
Rx	___ pill(s) every ___ hours as needed for _____				___ pill(s) every ___ hours as needed for _____			
DATE								

Medicine								
Rx	___ pill(s) every ___ hours as needed for _____				___ pill(s) every ___ hours as needed for _____			
DATE								

Surgeon's Office Number: _____

Knee Replacement Pre-Op Instructions Checklist

Your surgeon's office has scheduled you for Knee Replacement Surgery here at UnaSource Surgery Center. About 1-2 weeks prior to your scheduled surgery date one of our UnaSource team members will call to review your health history, provide instructions for surgery, and answer your questions.

You will need a responsible adult to accompany you to the surgery center, this person must be 18 years of age or older and must stay at the facility the entire time you are here. We allow one visitor per person; 2 visitors are allowed if the patient is under 18 years of age. Once discharged, a responsible adult must stay with you 24-48 hours after surgery.

Be sure to complete these 2 important steps prior to our call:

- Watch our educational Joint Replacement Video here: www.unasourcesurgery.com/joint-replacement/ Note: there is NO surgical footage in this video
- Flip through this Joint Replacement Experience Guide, and use the notes page to jot down your questions.

Our joint replacement video and experience guide are wonderful resources to prepare you for a successful surgery and recovery.

Day Before Surgery

- You will receive your arrival time the day before your surgery between the hours of 3pm-5pm. If you are scheduled on a Monday, you will receive a call the Friday before your surgery.
- **Follow all instructions given to you regarding which of your medications to take and which of your medications to hold.**
- Do not eat anything after midnight the night before your surgery. You may have clear liquids (any liquid you can see through) up to three hours before your arrival time.
- **Drink 10-12 ounces of a carb rich clear drink 3 hours prior to arrival or the night before** if your arrival is early. **Please choose between 100% no sugar added white grape juice, Gatorade, or Ensure Pre-Surgery Clear.**
- Please shower with Chlorhexidine Gluconate 4% (CHG) skin cleanser 2 days before and the morning of surgery. Do not apply any creams or lotions and do not shave the surgical site.

Diabetic or Weight Loss Medications

- If you are diabetic and you test your blood sugar regularly, please check it within **two hours** of your arrival time and let our staff know your result.
- If you take any of the following medications Ozempic, Wegovy, Trulicity, Mounjaro, Adlyxin, Byetta, Zepbound, Rybelsus, Victoza, or any other injectable type II diabetic or weight loss medications, **please hold that medication for 1 week. You are allowed clear liquids only for 24 hours prior to your day of surgery.**

Day of Surgery

- Wear loose fitting, comfortable clothes. Shoes should be supportive with closed heels.
- Remove all jewelry and body piercings. Leave all personal belongings at home.
- Remove contact lenses, bring glasses if applicable.

What to Bring

- Bring your insurance card, picture ID, and any expected payments.
- Bring a brace if prescribed for your recovery.
- If the nurse instructed you of the need to provide a urine sample upon arrival, please let our staff know when you arrive.




Notes:

Notes:

Notes:

Joint Replacement Education Video

Watch online at:
www.unasourcesurgery.com/joint-replacement



UnaSource Surgery Center

4550 Investment Drive, Suite 100

Troy, MI 48098

Phone: 248.265.4600

www.unasourcesurgery.com

www.facebook.com/unasourcesurgery